Mepilex[®] Border Flex

Oval shaped all-in-one flexible foam dressing



- Multilayered absorbent foam with flex technology
- Minimizes pain and trauma at dressing change^{2,3,5}
- Advanced fluid handling promotes optimal moisture balance and minimizes the risk of maceration^{4,6}
- Borders designed to adapt to body contours¹

Proven choice for a better outcome

Safetac*, pioneered by Mölnlycke, delivers above and beyond the ordinary. Proven to help optimize the wound healing journey and even prevent wounds, dressings with Safetac are the safe choice for patients and a champion for higher standards in wound care.

* A unique proprietary technology exclusive to Mölnlycke Health Care

In fact, we have a wealth of evidence that supports the clinical and economic benefits of dressings with Safetac, including Mepilex®, Mepitel®, Mepiform® and Mepitac®. To date, these dressings have helped millions of patients worldwide⁷⁻⁹.





Mepilex[®] Border Flex

Directions for use



1. Gently cleanse the wound; dry surrounding skin. Choose a size that will allow the dressing to overlap the wound area by at least 2 cm. Remove the release film.



2. Lay absorbent foam pad over wound area. Gently smooth adherent border in place. Do not stretch. Mepilex® Border Flex can remain in place for up to 7 days depending on drainage and wound condition.



How Mepilex[®] Border Flex works

Mepilex[®] Border Flex is an all-in-one oval foam dressing with advanced fluid handling capabilities, dynamic permeability⁴ and all the advantages of Safetac[®] soft silicone technology. The shapes of its borders have been designed to adapt to the body contours and to distribute forces on the borders to minimize rolling and increase both conformability and ability to stay in place. It is waterproof, viral proof^{*} and bacterial proof^{*} with a multilayered wound pad construction designed to promote optimal moisture balance and fluid uptake.^{34,6}

Benefits of Mepilex® Border Flex

- Highly conformable¹ to adapt to body contours
- Designed for enhanced ability to stay in place
- Minimizes pain and trauma at dressing changes^{2,3,5}
- Seals wound margins and reduces risk of maceration^{4,6}
- Can remain in place for several days depending on the condition of the wound
- Can be lifted and adjusted without losing its adherent properties
- Shower proof

Mepilex® Border Flex Assortment (Sterile packed)

Art. no	Size cm	Pcs/Box	Pcs/Case	
283570	7.8 x 10	5	50	
283300	13 x 16	5	35	
283400	15 x 19	5	45	

*Notice: For Mölnlycke licensed product details including indications and precautions, please refer to www.molnlycke.ca



References:

* microbes larger than 25 nm

 Tensile force, Conformability, in vitro tests. Mölnlycke Health Care lab report. 201303 01-002. 2. Upton D. et al. The Impact of Atraumatic Vs Conventional Dressings on Pain and Stress in Patients with Chronic Wounds. Journal of Wound Care, 2012. 3. White R. et al. Evidence for atraumatic soft silicone wound dressing use. Wounds UK, 2005. 4. Fluid handling capacity, in vitro tests. SMTL lab report SMTL 13/416/1.
S. White R. A. Wultinational Survey of the assessment of pain when removing dressings. Wounds UK, 2008. 6. Wiberg A.B. et al. Preventing maceration with a soft silicone dressing: in-vitro evaluations. Poster presented at the 3rd Congress of the WUWHS, Toronto, Canada, 2008.

Mölnlycke 1-800-494-5134 www.molnlycke.ca

Mölnlycke Health Care AB, Box 13080, SE-40252 Göteborg, Sweden. Phone +46317223000. The Mölnlycke Health Care, Mepilex Border® Flex, and Safetac® names and respective logos are registered globally to one or more of the Mölnlycke Health Care Group of Companies. © 2019 Mölnlycke Health Care AB. All rights reserved.

Indications*

Mepilex[®] Border Flex is designed for a wide range of exuding wounds such as pressure ulcers, leg and foot ulcers and traumatic wounds e.g. skin tears and surgical wounds.

Mepilex[®] Border Flex can also be used on dry/necrotic wounds in combination with gels.

Mepilex[®] Border Flex may be used as part of a prophylactic therapy to help prevent skin damage, e.g. pressure ulcers, postoperative blistering.

Precautions*

- Do not use on known sensitivity to the dressing or its components. In the case of signs of clinical infection, consult a health care professional for adequate infection treatment.
- Do not use Mepilex[®] Border Flex together with oxidising agents such as hypochlorite solutions or hydrogen peroxide.
- The use of dressings as part of a prophylactic therapy does not preclude the need to continue to develop and follow a comprehensive pressure injury prevention protocol, i.e. support surfaces, positioning, nutrition, hydration, skin care and mobility.

